

# Factsheet on preterm birth in Ghana

## What is preterm birth?

A preterm birth is when a baby is born alive but before 37 weeks of pregnancy are completed.<sup>1</sup> In Ghana, one in three newborn deaths are due to preterm birth complications (2013 estimate).<sup>2,a</sup> Babies born too soon are approximately seven times more likely to die during the first four weeks of their lives than babies born at term.<sup>3</sup>

In addition, preterm survivors often suffer from lifelong disabilities, such as visual and hearing impairments, chronic lung disease, long term cardiovascular ill-health, learning and behavioural impairments.<sup>1</sup> Prematurity has a far-reaching impact on their development and on their health as children and adults.

## Preterm birth is a public health problem in Ghana

Ghana ranks 23<sup>rd</sup> in the world for the number of preterm births.<sup>4,5,b</sup> Every year, 128,000 babies are born before 37 weeks of pregnancy are completed.<sup>4,5,b</sup> This is one in every 7 babies born alive.<sup>4,c</sup> Prematurity is the leading cause of death for babies in the first month of life<sup>2</sup>, with 7,900 newborn deaths due to preterm birth every year in Ghana<sup>2,6,d</sup>.

## What can be done in Ghana?

Many of these deaths could be prevented, through both prevention of preterm birth, and care for the premature baby.

- Ⓔ Adolescent pregnancy, older age pregnancy, short time gaps between births, chronic and infectious diseases, and unhealthy pre-pregnancy weight increase the risk of preterm birth<sup>1</sup>
- Ⓔ Family planning strategies, including birth spacing and adolescent-friendly services, can reduce the risk of preterm birth<sup>1</sup>
- Ⓔ Encourage early initiation and exclusive breastfeeding<sup>1</sup>
- Ⓔ Quality antenatal and postnatal care, along with skilled care at delivery, can help to reduce deaths due to preterm birth. Investing in trained and equipped health workers is key, with a focus on the 48 hours surrounding birth<sup>1</sup>
- Ⓔ In August 2015, the World Health Organization (WHO) published new recommendations to improve the survival and health of babies born too soon<sup>7</sup>. Decision-makers should review and revise national clinical guidelines and essential medicines lists; promote the new recommendations widely; strengthen the skills of health workers to deliver the interventions; support health workers to accurately assess gestational age; and monitor adherence to the guidelines.<sup>7</sup>

In 2014 Ghana committed to the Every Newborn Action Plan<sup>8</sup> and launched a national Newborn Health Strategy and Action Plan to speed up the reduction of newborn deaths. These strategies set out the priority solutions, and call for a united effort to dramatically maternal and newborn deaths, and preventable stillbirths. We must prioritise the evidence-based, cost-effective and feasible solutions in all health facilities that provide pregnancy and delivery services, as well as ensure better data collection on pregnancy outcomes to enable responsive action where it is most needed.

**Now is the time to scale up quality care to all women and newborns for them to survive, thrive and transform society!**





## Prematurity in Ghana

|                                                                                            |                |
|--------------------------------------------------------------------------------------------|----------------|
| <b>Number of newborn deaths in the first month of life (2015)<sup>6</sup></b>              | <b>25,400</b>  |
| Mortality rate in the first month of life (per 1000 babies born alive) (2015) <sup>6</sup> | 28/1000        |
| <b>Preterm birth rate (% of babies born alive) (2010)<sup>4</sup></b>                      | <b>14.5 %</b>  |
| Country rank for preterm birth rate (2010) <sup>4</sup>                                    | 14             |
| <b>Number of babies born preterm (2015)<sup>4,5,b</sup></b>                                | <b>128,000</b> |
| Country rank for number of babies born preterm (2015) <sup>4,5,b</sup>                     | 23             |
| <b>Number of newborn deaths due to preterm birth complications (2015)<sup>2,6,d</sup></b>  | <b>7,900</b>   |
| Share of newborn deaths caused by preterm birth complications (2013) <sup>2,a</sup>        | 31%            |
| <b>Demand for family planning satisfied (2014)<sup>9</sup></b>                             | <b>47%</b>     |
| Antenatal care of four or more visits (2014) <sup>9</sup>                                  | 87%            |
| <b>Skilled attendant at delivery (2014)<sup>9</sup></b>                                    | <b>74%</b>     |
| Newborns who did not receive any postnatal care visit after birth (2014) <sup>9</sup>      | 70%            |
| <b>Women initiating breastfeeding within 1 hour of birth (2014)<sup>9</sup></b>            | <b>56%</b>     |
| Proportion of babies aged 0-5 months who are exclusively breastfed (2014) <sup>9</sup>     | 52%            |

## References:

1. March of Dimes, The Partnership for Maternal Newborn & Child Health, Save the Children, World Health Organization. (2012). *Born Too Soon: The Global Action Report on Preterm Birth*. (Eds) Howson, C.P., Kinney, M.V., & Lawn, J.L. Geneva: WHO.
2. Liu, L., Oza, S., Hogan, D., Perin, J., Rudan, I., Lawn, J. E., Cousens, S., Mathers, C., & Black, R. E. (2014). Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. Supplementary appendix. *The Lancet*, 6736(14), 430-440.
3. Katz, J., Lee, A. C. C., Kozuki, N., Lawn, J. E., Cousens, S., Blencowe, H., Ezzati, M., Bhutta, Z., Marchant, T., Willey, B., Adair, L., Barros, F., Baqui, A., Christian, P., Fawzi, W., Gonzalez, R., Humphrey, J., Huybregts, L., Kolsteren, P., Mongkolkeha, A., Mullany, L., Ndyomugenyi, R., Nien, J., Osrin, D., Roberfroid, D., Sania, A., Schmiedel, C., Silveira, M., Tielsch, J., Vaidya, A., Velaphi, S., Victora, C. G., Watson-Jones, D., & Black, R. E. (2013). Mortality risk in preterm and small-for-gestational-age infants in low-income and middle-income countries: a pooled country analysis. *The Lancet*, 38(9890), 417–25.
4. Blencowe, H., Cousens, S., Oestergaard, M. Z., Chou, D., Moller, A.-B., Narwal, R., Adler, A., Garcia, C.V., Rhode, S., Say, L., & Lawn, J. E. (2012). National, regional, and worldwide estimates of preterm birth rates in 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. Supplementary appendix. *The Lancet*, 379(9832), 2162–2172.
5. United Nations, Department of Economic and Social Affairs, Population Division (2015). World Population Prospects: The 2015 Revision. File INT/1: Interpolated demographic indicators by major area, region and country, annually for 1950-2100. Retrieved on November 2, 2015, from <http://esa.un.org/unpd/wpp/DVD/>
6. UN Inter-agency Group for Child Mortality Estimation (IGME). (2015). Under-five mortality rate, Infant mortality rate, neonatal mortality rate, and number of deaths. Retrieved November 4, 2015, from <http://www.childmortality.org/>
7. World Health Organization. (2015). WHO Recommendations on Interventions to Improve Preterm Birth Outcomes. Geneva: WHO.
8. World Health Organization, & UNICEF. (2014). Every Newborn: An Action Plan to End Preventable Deaths. Geneva: WHO.
9. Ghana Statistical Service, Ghana Health Service, and ICF International. (2015). Ghana Demographic and Health Survey 2014. Rockville: GSS, GHS, & ICF International.

## Notes:

- a. In Ghana the estimated number of newborn deaths for 2013 is 23,110 and the number of deaths due to preterm birth complications is 7,159.<sup>2</sup> These estimations were used to calculate the proportion of neonatal deaths due to preterm birth complications:  $(7,159/23,110) \times 100 = 30.978\%$  (1 in 3).
- b. We calculated the number of preterm births in Ghana by using the most recent preterm birth rate (2010)<sup>4</sup> and applying this to the 2015 estimate of live births<sup>5</sup>. In Ghana, the estimated number of live births for 2015 is 883,700<sup>5</sup>. The 2010 preterm birth rate is applied to the 2015 live birth estimate:  $(14.48906/100) \times 883,700 = 128,039.82322$  (rounded to 128,000). Countries are ranked by number of preterm births based on these estimates for 2015.
- c. The preterm birth rate is 14.49%<sup>4</sup>; this is equivalent to 1 baby out of every 7 being born preterm.
- d. We calculated the number of deaths due to preterm birth complications in the first 28 days of life in Ghana by using the latest data (2013) on the proportion of newborn deaths from preterm birth (30.978%)<sup>2,a</sup> and applying this to the 2015 estimate of the number of newborn deaths (25,402)<sup>6</sup>. This is equivalent to:  $(30.978/100) \times 25,402 = 7869$  (rounded to 7,900). 7,900 is the estimated number of deaths due to preterm birth complications in the first 28 days of life in Ghana in 2015. Including those who die from preterm birth complications between 0 and 5 years the figure is 8,400 (estimate provided by Healthy Newborn Network, 2015).

This factsheet was produced in November 2014 and updated in November 2015.